

Carbon Management



ISSN: 1758-3004 (Print) 1758-3012 (Online) Journal homepage: https://www.tandfonline.com/loi/tcmt20

Climate-smart land use requires local solutions, transdisciplinary research, policy coherence and transparency

Sarah Carter, Bas Arts, Ken E. Giller, Cinthia Soto Golcher, Kasper Kok, Jessica de Koning, Meine van Noordwijk, Pytrik Reidsma, Mariana C. Rufino, Giulia Salvini, Louis Verchot, Eva Wollenberg & Martin Herold

To cite this article: Sarah Carter, Bas Arts, Ken E. Giller, Cinthia Soto Golcher, Kasper Kok, Jessica de Koning, Meine van Noordwijk, Pytrik Reidsma, Mariana C. Rufino, Giulia Salvini, Louis Verchot, Eva Wollenberg & Martin Herold (2018) Climate-smart land use requires local solutions, transdisciplinary research, policy coherence and transparency, Carbon Management, 9:3, 291-301, DOI: 10.1080/17583004.2018.1457907

To link to this article: https://doi.org/10.1080/17583004.2018.1457907





OPEN ACCESS (Check for updates



Climate-smart land use requires local solutions, transdisciplinary research, policy coherence and transparency

Sarah Carter ^[ba,b], Bas Arts^c, Ken E. Giller^d, Cinthia Soto Golcher^c, Kasper Kok ^[be], Jessica de Koning^{c,f}, Meine van Noordwijk^{d,g}, Pytrik Reidsma^d, Mariana C. Rufino^h, Giulia Salvini^f, Louis Verchotⁱ, Eva Wollenberg^j and Martin Herold (10)

^aWageningen University & Research, Laboratory of Geo-Information Science and Remote Sensing, Droevendaalsesteeg 3, Wageningen, 6708 PB Netherlands; ⁵Center for International Forestry Research, Bogor Barat, 16115 Indonesia; ^cWageningen University & Research, Forest and Nature Conservation Policy Group, Wageningen, Netherlands; dWageningen University & Research, Plant Production Systems, Wageningen, Netherlands; eWageningen University & Research, Soil Geography and Landscape Group, Wageningen, Netherlands; Wageningen University & Research, Laboratory of Geo-Information Science and Remote Sensing, Wageningen, Netherlands; Wageningen, Wageningen, Netherlands; Wageningen, Wag Agroforestry Centre (ICRAF), Bogor, Indonesia; hCenter for International Forestry Research, Nairobi, Kenya; Center for International Forestry Research, Bogor Barat, Indonesia; ¹Gund Institute, University of Vermont and CGIAR Research Program on Climate Change, Agriculture and Food Security, Burlington, United States; Wageningen University, Laboratory of Geo-Information Science and Remote Sensing, Wageningen, Netherlands

ABSTRACT

Successfully meeting the mitigation and adaptation targets of the Paris Climate Agreement (PA) will depend on strengthening the ties between forests and agriculture. Climate-smart land use can be achieved by integrating climate-smart agriculture (CSA) and REDD+. The focus on agriculture for food security within a changing climate, and on forests for climate change mitigation and adaptation, can be achieved simultaneously with a transformational change in the land-use sector. Striving for both independently will lead to competition for land, inefficiencies in monitoring and conflicting agendas. Practical solutions exist for specific contexts that can lead to increased agricultural output and forest protection. Landscape-level emissions accounting can be used to identify these practices. Transdisciplinary research agendas can identify and prioritize solutions and targets for integrated mitigation and adaptation interventions. Policy coherence must be achieved at a number of levels, from international to local, to avoid conflicting incentives. Transparency must lastly be integrated, through collaborative design of projects, and open data and methods. Climate-smart land use requires all these elements, and will increase the likelihood of successful REDD+ and CSA interventions. This will support the PA as well as other initiatives as part of the Sustainable Development Goals.

KEYWORDS

Climate policy; climate-smart agriculture: deforestation: food security; REDD+

Introduction

The ambitious goals set at the 21st conference of parties (COP) of the United Nations Framework Convention on Climate Change (UNFCCC) in Paris have since been discussed in Bonn and Marrakesh. The Road Map for Global Climate Action notes a long-term move in focus from mitigation action toward adaptation [1], which will also be reflected in funding allocation [2]. The Marrakesh Action proclamation promises actions in the agriculture sector, primarily ensuring food security and enhancing the ability to deal with climate change impacts on agriculture [3]. There was further progress on agriculture in Bonn; the links between agriculture and climate change were included as a discussion point, with options such as increasing soil carbon mentioned [4]. At the same time, the Paris Climate Agreement (PA) features forest-based mitigation as a key mitigation strategy as well as working to secure food production. This presents a potential conflict between forests and agriculture.

The land-use sector is unique in its large potential for negative emissions, besides climate engineering options such as carbon capture and storage [see 5], and therefore must be fully utilized. The focus on food security and food production in the PA must leverage the potential synergies between adaptation and mitigation, which will support forest protection [6]. A crucial point of entry relates to linking existing concepts that address climate change mitigation and adaptation in the land-use sector, such as climate-smart agriculture (CSA) and REDD+. CSA aims to tackle three main objectives: sustainably increasing agricultural productivity and incomes; adapting and building resilience to climate change; and reducing and/or removing greenhouse gas emissions, where possible [7]. REDD+, a

forest-based mitigation mechanism, aims to reduce emissions from deforestation and forest degradation, and to conserve forests and enhance forest carbon stocks by reducing pressure from drivers of deforestation and by managing forests sustainably. The agricultural drivers of deforestation, and their links to food security, should also be well understood. Commercial agriculture, which is often the result of large-scale land acquisitions, can have a large impact on forests [8], and can lead to food insecurity for smallholders [9]. Smallholder farming, in contrast, has a large role to play in terms of food security [10]. These key differences must be addressed when planning interventions.

The primacy of food production is clear in global strategies, not only in the PA (and the main text of the UNFCCC) but also in the Sustainable Development Goals [11]. With the increasing global population, this focus is crucial, as population increases are not matched by food production increases [12], and in Africa, population growth is expected to be the dominant driver of food insecurity, above climate change [13]. As such, the focus of CSA leans toward food security and adaptation, with mitigation being pursued only 'where possible'; a shift from earlier definitions in which it was described as a mandatory component [14]. In addition, the mitigation potential of agriculture and forests varies greatly from country to country [15], as well as the motivation within a country to focus on one or the other of the aims [16]. A major challenge in reconciling the two is the different accounting bases for food production relative to consumer demand, and area-based carbon balance in land use. For this reason, this paper focuses on the forest-sparing potential of interventions in the agricultural sector that seek to increase food production, one element of food security [17].

The starting point for the analysis is the need to reconcile competing claims on land for climate mitigation (including forest protection and biofuel production) and for food production. Since deforestation is closely linked with agriculture [18,19], REDD+ will fail if agricultureal expansion into forests persists [20,21]. A review of REDD+ readiness documents reveals that most forest-related policies tend to overlook agricultural drivers of deforestation [22]. Initiatives that increase production on existing agricultural land can, in a planned economy, reduce the need to expand agriculture into, for example, high-carbon landscapes (forests, woodlands, wetlands). In a market economy, however, the land-sparing effects of intensification may only be achieved if it induces commodity prices to drop to a level where expansion and less-intensive production become uneconomic uses of land and labor; otherwise, increasing profits from yield improvements can lead to increased land acquisition for agriculture. This can, however, be avoided, if agricultural intensification (undertaken with CSA principles) is coupled with forest protection mechanisms (e.g. policies or interventions that directly address the agriculture drivers) to ensure that forest-sparing actually occurs [23,15,24–26]. While REDD+ may operate with area-based ('simple') scaling rules, global markets for agricultural commodities imply more complex scale relationships for CSA, and hence for the way REDD+ and CSA are related to each other [27]. These feedbacks at different scales call for an exploration of how REDD+ and CSA could be linked, and what the mutual synergies and trade-offs might be. This article proposes an approach in which not all agricultural expansion is avoided, but in which emissions from agriculture-driven deforestation are minimized, and food security is protected.

Support to develop solutions that reduce deforestation may be found in the emerging science on the dependence of hydroclimates on vegetative cover [28], which further highlights the links between CSA and REDD+. For example, those interventions to secure food production and those to protect forests can be mutually beneficial: forests provide numerous ecosystem services that can increase the adaptive capacity of agriculture, which reduces vulnerability to climate change [29,28]. Forest-generated humidity, soil stability and soil fertility maintenance are crucial services for agriculture, but also support communities by providing drinking water and reducing risks such as landslides and flooding. Additionally, forest products provide safety nets for local communities when agricultural yields decline because of climate change impacts [30], further supporting the goal of securing food production [31,32].

A roadmap for transformational change

Currently, there is a lack of coherent policies, and conflicting incentives, in the agriculture and forest sectors, implying that emissions reductions can only be realized through a deep transformational change. Transformational change is a move away from business as usual and is likely to include a shift in practices, a change in commodities, innovative policies, and/or financing actions [33]. The authors envisage that changes can be applied incrementally or stepwise; however, major changes across sectors and at all levels will be required [33,34].

To address the challenge of competition for land, while promoting forest-based mitigation efforts without compromising food production, this paper proposes an area-based integration of REDD+ and CSA at intermediate (e.g. local government or 'jurisdictional') scales, to enhance synergies and reduce trade-offs. The authors argue that the costs of not integrating will lead to competition for land and other resources, inefficiencies in monitoring and conflicting agendas, which leads to less success in reducing forest loss and in protecting food security. An integrated approach requires:

- Practical local solutions that integrate CSA and REDD+ and reduce emissions at the landscape scale;
- Transdisciplinary research approaches and priorities, in the agricultural, forest and social sciences and in non-scientific communities;
- Policy coherence at (inter)national and local levels;
- Transparency in reporting and engagement.

Practical local solutions

The definition of what is and what is not CSA is important if economic incentives and policy recognition matter. This article proposes that practices should only be classed as CSA when they, in addition to enhancing food security and increasing resilience, achieve reductions in greenhouse gas emissions (mitigation) at the landscape level (including, importantly, the life cycle of the product), and avoid expansion of agriculture into forests, which should be avoided where possible. Sa et al. [35] used a similar definition for activities that mitigate climate change and advance food security in South America. They stated these activities must have 'low carbon dioxide (CO₂) emissions from land use (LU) and land use change (LUC) in response to agricultural best management practices' [35]. Only activities that meet these criteria were considered suitable by the authors to contribute to the low-carbon agriculture (LCA) strategy as part of Brazil's program on

low-carbon agriculture [36]. The activities identified were diverse, and included restoration of degraded pasture, biological N fixation and plantations of commercial forests. In all the scenarios assessed by the study, food production increased. In order for this to occur, the expansion of agricultural land was required. In these cases, emissions from the expansion event should be estimated and used to guide decision-making as to whether and where agriculture should expand.

Calculating the emissions intensity of crops is a useful tool to understand the impact of production and also to set targets for mitigation [37]. Different outcomes in terms of the best options for mitigation can be found using different accounting methods [38], so using a number of approaches and comparing results can be useful for decision makers. The 'carbon debt' when establishing oil palm plantations in Indonesia has been discussed by van Noordwijk et al. (Figure 1) [38]. In this case, an optimal fertilizer level is determined by the production levels at which net emission savings per unit of biofuel are maximized. These figures are then compared to the carbon debt of clearing land - including draining of peatlands, and the loss of forests. The carbon debt and emissions from fertilizer were the dominant factors in the whole life-cycle assessment - showing how important the impacts on the forest sector are for mitigation efforts in agriculture. Palm oil produced on peat soils, and in some cases mineral soils, was found not to meet the current




Figure 1. Information flow in an assessment of the emission footprint per unit palm oil, and a subsequent step to estimate the percentage emission saving in biofuel use (adapted from van Noordwijk et al. [38]). FFB: Fresh fruit bunches; CPO: Crude palm oil.

EU emissions threshold for biofuel production, which requires at least 35% emissions savings compared to fossil fuels.

In van Noordwijk et al. [38], an optimum fertilizer application rate can be identified for a given relationship between fertilizer and yield. Attention to other management choices can, however, shift the relationship between fertilizer and yield, and current fertilizer use on smallholder oil palm farms can be excessive and inefficient [39].

Similarly, Sa et al. [35] assessed emissions associated with land-use change over time using the concept of ecosystem carbon payback time (ECPT). ECPT considers how long the annual savings from the LCA activities (in comparison with non-LCA) can repay the debt of conversion (e.g. clearing the forest in the conversion to agriculture). ECPT, expressed in years, represents the time that the intervention has to be effective in order to offset the land use and land use change emissions from the intervention. Implementing activities in areas with the lowest ECPT is a way of minimizing the impact, and the restoration of degraded pastures was found to be one of the most advantageous scenarios, although it still had an ECPT of 56 to 188 years [35]. In the case of Sa et al. restoration of pastures was achieved by application of fertilizers, lime and gypsum (accounting for associated emissions related to their application), as well as altering species composition and introducing forage management to increase the carrying capacity. The restoration of grazing land has also been implemented as a National Appropriate Mitigation Action (NAMA), and has been estimated to fulfil 10–12% of pledged emission reductions for the year 2020, due to land spared from deforestation [40]. Restoration of pastures can be a promising tool to reduce agricultural expansion into forests (see e.g. [40], although pasture activity has been found to have mixed impacts on deforestation [41]), and should therefore be considered a priority mitigation action in the agriculture sector. However, in many cases, degraded land has marginal potential for production, even after costly rehabilitation. Lal [42] suggested that agriculturally marginal and degraded soils should be avoided for this purpose (and could be set aside for nature), and that lands with potential for higher yields could be restored, for example those in areas of sub-Saharan Africa and South Asia where there is potential for tripling or quadrupling of yields. Over 1.3 billion people, however, use degraded agricultural land, particularly in the drylands [43], and in general larger soil C sequestration rates can be achieved from interventions to increase C sequestration in degraded soils than from other soils [44]. There are a number of GHG mitigating practices that can be implemented to restore degraded and marginal land, such as the use of cover crops, avoiding overgrazing, using crop residues and organic composts and agroforestry; however, employing these is not always straightforward [45]. The conditions under which yields in sub-Saharan Africa and South Asia could be increased while sparing land should be investigated, utilizing the existing body of evidence on successful CSA interventions [7; see also e.g. 35,38,39,42]. The dichotomy between forest and agriculture itself may prevent the emergence of optimal solutions, as for example in a landscape where fodder for dairy cows can be derived from any part of the landscape [46].

The sink capacity of forests can provide a large and cost-effective contribution toward climate change mitigation [47]. However, agricultural systems can also provide mitigation benefits, as well as contributing to forest sparing. In fact, a recent global study of trees in agricultural lands suggested a considerable carbon stock that has so far been ignored in the accounting [48]. Agroforestry and soils represent a sink potential and technologies such as biogas digesters a promising mitigation potential, and measures related to these potentials have been implemented without compromising agricultural production [49]. Agroforestry has one of the highest land-based potentials to sequester carbon [50,21,51], and provides the adaptation benefits aimed at by CSA. It is also reversible, however, so land can be converted back to treeless agriculture should local priorities change. Reducing emissions from soils storage is possible through a number of management interventions such as increasing inputs that contain C, avoiding over-fertilization of crops, and altering irrigation and tillage regimes [44]. This has received recent attention, for example through the Soils for Food Security and Climate 4% Initiative [42,52]. Knowledge gaps exist, however, for example in soil C sequestration processes, turnover and stabilization [45], as do potential trade-offs between soil C sequestration and N2O emissions [53], making it controversial not least because it is easily reversible. There is currently also an active debate on the potential of the 4-per-mille initiative, and whether the claims (of offsetting 30% of global GHG emissions [44]) are feasible. Although the difficulties of engaging large numbers of people who are using agricultural lands around the world, and the technical challenges in monitoring and verifying results, are discussed [44], other points such as the feasibility of including currently unmanaged rangelands, time to reach carbon equilibrium and depth at which soils can be managed are still being debated [54, among others]. However, identifying areas with the potential to increase soil C stocks at a rate of 4% could contribute to climate mitigation efforts, and soils remain an important consideration in the context of this paper, in their role in forestsparing and food security.

Although interventions exist that not only deliver increased production within a changing climate (food security), but also avoid land-use change (particularly

from forests or other high carbon landscapes), robust and inclusive methods to identify which ones can provide the largest mitigation benefits must be used.

Transdisciplinary research approaches and priorities

Transdisciplinary research is vital to set priorities for climate change actions in the land-use sector. Research should recognize the varied interests of each actor by answering questions and delivering results that stakeholders find useful [55]. Approaches should account for the disconnect between the timescales at which various actors operate - for example, climate science considers longer timescales than those at which farmers, and also foresters, make decisions. Using backcasting to develop transition pathways is one way to evaluate options to achieve multiple sustainability objectives [e.g. 5]. In addition, models and tools should be able to assess the outcomes of various interventions related to adaptation, mitigation and food production [56-58]. Trade-offs for stakeholders are inevitable, and need to be evaluated (e.g. in the context of the landscape approach). Smallholder farmers are often key stakeholders who should be consulted, and who will then assess whether they are willing to engage in such interventions. Engel and Muller [59] discuss the potential for incorporating CSA into a payments for ecosystem services (PES) context (e.g. REDD+), and highlight the potential for PES to address reasons why some farmers do not adopt CSA. A PES approach was found to deal with issues such as an unwillingness to wait for medium- or long-term productivity and an aversion to risk, as short-term costs are often required [59]. Other barriers identified included lack of information, insecure tenure and weak property rights. These issues can be addressed by incorporating CSA into the framework of REDD+ projects. Securing tenure and community

engagement are key processes for REDD+ implementation.

Landscape approaches could provide a platform for assessing the benefits and challenges of integrating mitigation and adaptation [27,60,61] (also for emissions accounting, as discussed previously). To involve and improve communication between stakeholders and researchers in the landscape, methods using both quantitative and qualitative data can be used (e.g. role playing games [62,63], and innovative participatory scenario planning [64]). Increasing availability of remote sensing data, emerging technologies and community-based monitoring can be the focus of research streams aiming to monitor results from both CSA and REDD+ together [65,66]. Measuring the impacts of ongoing initiatives is key to understanding how to implement a landscape approach.

To have the greatest impact, interventions in the agricultural sector that promote forest land-sparing should be implemented first in areas where most agriculture-driven deforestation and emissions from agriculture are occurring [67]. These were identified at the national level by Carter et al. [15]. Emission hotspots have been identified at a much higher spatial resolution by Roman-Cuesta et al. [68], through a collaborative effort from the forest and agricultural communities. Hotspots covering 25% of the tropical area are responsible for 70% of the tropical agriculture, forestry and other land use (AFOLU) emissions. All continents have hotspots, and they cover a variety of biomes across the tropics - so these make promising locations for interventions [68]. Typically emissions in hotspots are dominated by forest emissions (69%; e.g. fire, deforestation and wood harvesting), highlighting the important role of forests in mitigation, although livestock and paddy rice also produce high emissions (Figure 2).

One important point for research priorities from the example of Roman-Cuesta et al. [68] is that hotspot




Figure 2. Contribution of the leading emission sources (grouped into forests, crops and livestock) % per pixel (0.5°), 2000–2005. Forest emission sources include fire, deforestation and wood harvesting. Crop emissions include paddy rice, cropland soil and croplands over drained histosols. Livestock includes enteric fermentation and manure management emissions. Colors represent the strength of the emissions for the three sources (e.g. fuchsia in Asia represents equal emissions from livestock (red) and crops (blue). Dark areas are areas of low emissions. Adapted from Roman-Cuesta *et al.* 2016 [68].

regions have the highest uncertainties. These are up to 30% of the mean AFOLU emissions, so research needs to focus on reducing uncertainties for emissions estimates [68]. This is because uncertainty in these estimates leads to uncertain potential benefits from interventions. Forests have the highest uncertainties associated with their emissions, with their uncertainties accounting for 98% of the AFOLU emission uncertainty. This is due to the combined effect of uncertain areas and uncertain carbon densities, so future research could focus on quantifying these variables [68]. The contribution of forests as a carbon sink therefore requires further research, as the impact of the recovery of carbon stocks after wood harvesting and fire could be better understood [68].

Once locations for mitigation initiatives have been identified, the impact of the initiatives must be monitored. Measuring and evaluating mitigation options for smallholder farmers (who occupy most land holdings in developing countries) is difficult due to the lack of available emissions factors and the diversity of smallholder farming systems [e.g. 58]. The needs of the organization doing the measuring also differ in terms of accuracy required, budget and scale of measurements, so one monitoring, reporting and verification system will not suit all scenarios. However, protocols have been developed that offer options for measuring performance according not only to mitigation and adaptation goals, but also to food security and local livelihoods, and which can be adjusted to suit the needs of the project developer [69]. Research could focus on testing and improving such approaches, as well as incorporating monitoring of forests and agricultural interventions using the same system.

Policy coherence

At the international level, agriculture and forests are often addressed through their own international platforms such as the Committee on Food Security for agriculture and the United Nations Forum on Forests (UNFF) for forests, and even through different UNFCCC Subsidiary Body for Scientific and Technological Advice (SBSTA) negotiation tracks. There has been a call for an SBSTA negotiation track on agriculture [70,71], but a better option might be to establish an SBSTA negotiation track that could accommodate landscape issues (including forests and agriculture) in one body. In addition to the UNFCCC, other international interventions and their targets are relevant for both forests and agriculture, including the Sustainable Development Goals, the Soils for Food Security and Climate 4‰ Initiative, the Aichi targets under the UN convention on Biological Diversity, and the land degradation neutrality target under the UN Convention to Combat Desertification. As well as conflicting goals and trade-offs relating to

forest and agriculture objectives within these, conflicts among them should be investigated.

At the national level, strong policy coherence is also important [23], and conflicting incentives (e.g. subsidies and tax breaks) should be removed and trade-offs addressed to create a supportive environment for both adaptation and mitigation in the land-use sector. Utilizing the institutional structures of REDD+ (specifically in its national policies, which have been relatively rapidly developed [72]) and integrating CSA should be explored as priorities, particularly where there are clear links between the two (e.g. high-yield palm oil in Peru [73]). Inspiration might be found in India's agroforestry policy adopted in 2014, which opted for an 'agriculture plus forestry' concept in a level playing field: any policy applied to one should be applied to the other, relative to measurable performance. In this case, specific funds were allocated for the harmonization of the approaches [74].

The Nationally Determined Contributions (NDCs) implemented under the UNFCCC relate to different components of the land sector, and could be a starting point for an integrated national approach. In fact, agriculture and forestry are already prominent components of many NDCs, with most parties using forest management and reforestation in their mitigation measures [16], so including agricultural NDCs or linking them to forest mitigation and adaptation could be further developed in future. Indeed, forest-based mitigation could provide a quarter of the emission reductions planned in several NDCs [75].

As the demand for agricultural products is responsible for much deforestation [76], the role of the private sector (including international markets) is important in connecting REDD+ and CSA. Private-sector commitments include green public procurement policies, for example the Roundtable for Sustainable Palm Oil and Round Table on Responsible Soy. One promising example from the private sector is the zero-deforestation commitments that many companies are adopting, even though the choice of net or gross zero deforestation creates space for a disparity of outcomes [77]. Despite these policies being adopted by some companies, a report from Forest 500 reveals that there are still many companies with weak or no policies on deforestation [64].

How to link commitments from the private sector to REDD+ is a key question. Research shows that there is a potential for agricultural value chains to be further integrated into REDD+ and CSA strategies [78]. Examples include 'jurisdictional approaches to zero-deforestation commitments' (JA-ZDCs) that allow policies such as REDD+ to be linked with private-sector interventions that also seek to reduce deforestation. Zero-deforestation policies and REDD+ already exist in parallel, so this option allows their numerous synergies (in terms of social and environmental goals) to be brought



Figure 3. Intersection of three strategies to reduce deforestation which centre around jurisdictional approaches to zero-deforestation commitments (JA-ZDCs). Adapted from [80].

together [79]. These approaches encompass a number of stakeholders and spheres in which they operate, including jurisdictions of governments (national or subnational), private-sector actors, and also a landscape perspective [80] (Figure 3). Notable examples include the Forest, Farms and Finance initiative, whose goal is to link interventions that aim to increase agricultural productivity and improve smallholder livelihoods with interventions that reduce deforestation [81]. Monitoring of these integrated approaches may also be more efficient. The Earth Innovation Institute has proposed a Territorial Performance System, which includes integrated incentives, an online monitoring system and multi-stakeholder governance. Other jurisdictional approaches to zero-deforestation commitments have been led by the private sector, including Marks and Spencer, and Unilever who pledged to preferentially source production from jurisdictions that adhere to a number of criteria including that the jurisdiction has a strategy for reducing emissions from deforestation [82]. Success of integrated approaches has been seen in the Brazilian Amazon, for example, where deforestation has been halted due to a combination of efforts from both the private sector and jurisdictional government, and voluntary efforts in the landscapes [83].

In addition to supply-side policies, policies or efforts to address demand for different types of food are needed. Demand-side measures have been suggested to have a greater mitigation potential (1.5-15.6 Gt CO_2 -eq. yr⁻¹) than supply-side measures (1.5–4.3 Gt CO2-eq. yr⁻¹) [84]. They are, however, difficult to achieve and require great societal change and commitment. One example requiring social commitment is diet change. Since there is a strong relationship

between income and meat intake, future increases in meat consumption are expected. Limiting animal protein intake can save land [85,86]. Examples of mechanisms that can achieve societal change are Voluntary Sustainability Standards (VSS) that inform consumers about the sustainability of goods [87]. Examples are Forest Stewardship Council (timber), Marine Stewardship Council (fish) and UTZ (coffee). Although they originate from private and civic sectors that cooperate to 'green' global value chains, governments can - and do in various cases - actively endorse these through their own policies (the Netherlands being an example). An integrated framework for analyzing where and how standards and certification emerge and evolve in the case of tropical timber, coffee, cacao, palm oil and rubber is provided by Mithöfer et al. [88]. VSS with very high social and environmental requirements provide information to consumers. However, they represent a small segment of the market, so product-wide commitments even with lower requirements could also lead to a larger impact. The Carbon Disclosure Project is a platform where companies can voluntarily report their environmental impacts, and most of the largest companies in the world participate [89]. Like voluntary standards, this information allows investors and consumers to make informed choices; however, the system remains voluntary. Mandatory reporting could provide more comprehensive information on emissions and carbon management practices, although reporting of practices to reduce emissions does not necessarily lead to a reduction in emissions [90]. The authors recommend that any future reporting should indicate the emissions reductions made by the company as well as reporting on forest impacts, both direct and indirect, from the supply chain. This would be part of monitoring the three components of CSA.

Transparency

Transparency should be understood as a catalyst for action by providing open and consistent data, definitions, assumptions and methodologies for an assessment of the credibility and reliability of land-use-sector mitigation and adaptation activities in both developing and developed countries. Transparency and open data allow the sharing of information between the forest and agriculture sectors, thus allowing the goals to be evaluated together. Enhancing transparency is now a fundamental step to make the bottom-up nature of the PA work in practice, and increases accountability for the various stakeholders involved [91]. This development is supported by an increasing set of free and open data and methods, for example through Global Forest Watch [92], OpenForis [93] and Global Open Data for Agriculture and Nutrition (GODAN) [94]. On the national level, enhancing transparency for emissions accounting, including estimates for mitigation

action impacts (for example NDCs) [75], is already one of the reporting criteria for the IPCC GHG inventories (transparency, accuracy, completeness, comparability, consistency) [95]. Such national reporting will be increasingly compared to independent data sources as part of the technical reviews and UNFCCC global stocktakes. National and other independent data sources should become available for local stakeholders to underpin their mitigation and adaptation activities. This should target non-state actors in particular, such as land owners, farmers, local communities and the private sector, to stimulate them to participate in and achieve more climate-smart land-use decisions and actions. Transparency for the private-sector initiatives (e.g. zero-deforestation commitments) is key to ensure that consumers trust in results, and independent assessments or third-party verification is a requirement for the voluntary sustainability standards mentioned earlier. Transparency should be multi-dimensional and interactive, and cover all stakeholders, thus being a pathway to engagement. Open and transparent sharing of information is essential for collaborative design and interactive processes to involve communities, for example using the participatory methods described by Salvini et al. [62]. This has been shown to reduce the risk of failure in mitigation activities, for example in the case of the voluntary carbon market that led to a decrease in credit prices [96]. Thus, transparency

 Provide data, case studies and guidance, which can be used to identify emissions reductions related to CSA activities that reduce agriculture expansion into forests, and can therefore assist in decision-making priorities and reporting at the national level, as well as contributing to UNFCCCcoordinated technical assessment processes and upcoming global stock-take(s);

should become a key element in planning, implemen-

tation and reporting of activities related to reducing

the impact of agriculture on forests, with a particular

need to address the following issues:

- Underpin multi-stakeholder processes, involving both the agriculture and forest sectors, for streamlining pathways to achieve land use sector mitigation and adaptation on the national (i.e. NDCs) and landscape scales, as well as increasing food production;
- Facilitate participatory monitoring for tracking and impact assessment of land-use mitigation activities involving the private sector engaged in zero-deforestation, civil society and government agencies.

Conclusions

New local integrated solutions, transdisciplinary research approaches and priorities, policy coherence

and transparency can all be independently sought and achieved, but transformational change will only happen when they are pursued simultaneously. We recommend that this transformational change include several components. First, that landscape-level accounting is used to assess the mitigation potential for interventions in the agricultural sector, which not only deliver increased production within a changing climate (food security), but also avoid land-use change (particularly from forests or other high carbon landscapes) where possible. Second, interventions in 'hotspot' regions, where the most emissions are occurring are given priority. In order to limit global average temperature increase to 1.5°C, attention should first be given to those emissions hotspots, and all aspects of the land sector - particularly the potential of the agriculture and forest sectors - should be utilized. Research should focus on decreasing uncertainties in emissions estimates in the land sector, so that more accurate mitigation potentials can be calculated. Third, policies at all levels must be coherent, and must support both mitigation and adaptation. At the international level, landscape issues should be discussed together. At the national level, institutional structures from REDD+ can be used to support CSA interventions, and these can also integrate private-sector actors, such as through JA-ZDC. Demand-side measures are also achieved and can be furthered through voluntary private-sector standards, for example. Fourth, transparency can be used to build trust, and to encourage the engagement of stakeholders. Support from the international community, including scientists, through the development of open data and tools can ensure that these initiatives are successfully implemented. This transformational change will also support other platforms, for example Sustainable Development Goals 2, 13 and 15 in particular, which focus on hunger, climate change and sustainable management of natural resources.

Acknowledgements

This research is part of CIFOR's Global Comparative Study on REDD+ (www.cifor.org/gcs). The funding partners that have supported this research include the Norwegian Agency for Development Cooperation (Norad) [Grant no. QZA-16/0110 No. 1500551] and the International Climate Initiative (IKI) of the German Federal Ministry for the Environment, Nature Conservation, Building and Nuclear Safety (BMUB) [Grant no. KI II 7 – 42206-6/75], and the CGIAR Research Program on Forests, Trees and Agroforestry (CRP-FTA), and Climate Change Agriculture and Food Security (CCAFS), with financial support from the CGIAR Fund Donors.

Disclosure statement

No potential conflict of interest was reported by the authors.



Funding

This work was supported by the Direktoratet for Utviklingssamarbeid [grant number QZA-16/0110 No. 1500551]; International Climate Initiative of the German Federal Ministry for the Environment, Nature Conservation, Building and Nuclear Safety (BMUB) [grant number KI II 7 - 42206-6/75].

ORCID

Sarah Carter (b) http://orcid.org/0000-0002-1833-3239 Kasper Kok (b) http://orcid.org/0000-0002-6319-9227 Martin Herold (http://orcid.org/0000-0003-0246-6886)

References

- 1. Tubiana L, El Haite H. Road map for global climate action [Internet]. (2016). Available from: http://newsroom. unfccc.int/media/658505/high-level-champions-climateaction-roadmap.pdf
- 2. UNFCCC. Long-term climate finance, Agenda item 10(a), Proposal by the President, Draft decision -/CP.22 [Internet]. (2016). Available from: http://unfccc.int/%5Cncoo peration_and_support/financial_mechanism/items/ 2807.php
- 3. UNFCCC. Marrakech action proclamation for our climate and sustainable development [Internet]. (2016). Available https://unfccc.int/files/meetings/ marrakech_nov_2016/application/pdf/marra kech_action_proclamation.pdf
- 4. UNFCCC. (2017). Advance unedited version. Decision -/CP.23. Koronivia joint work on agriculture. https:// unfccc.int/files/meetings/bonn_nov_2017/application/ pdf/cp23_auv_agri.pdf.
- 5. van Vuuren DP, Kok M, Lucas PL et al. Pathways to achieve a set of ambitious global sustainability objectives by 2050: explorations using the IMAGE integrated assessment model. Technol. Forecast. Soc. Change 98, 303-323
- 6. Duguma LA, Minang PA, Van Noordwijk M. Climate change mitigation and adaptation in the land use sector: from complementarity to synergy. Environ. Manage. 54 (3), 420-432 (2014).
- 7. FAO. Climate-Smart Agriculture Sourcebook [Internet]. (2013). Available from: http://www.fao.org/docrep/018/ i3325e/i3325e.pdf
- 8. Carter S, Manceur AM, Seppelt R, Hermans-neumann K, Herold M, Verchot L. Large scale land acquisitions and REDD+: a synthesis of conflicts and opportunities. Environ. Res. Lett. 12(3), 35010 (2017).
- 9. Cotula L. The international political economy of the global land rush: a critical appraisal of trends, scale, geography and drivers. J. Peasant Stud. . 39(3-4), 649-680 (2012).
- 10. FAO. The State of Food and Agriculture: innovation in family farming. (2014). Available from: http://www.fao. org/3/a-i4040e.pdf
- 11. UN. UN Sustainable Development Goals [Internet]. (2016). Available from: https://sustainabledevelopment.un.org/? menu=1300
- 12. Alexandratos N, Bruinsma J. World Agriculture Towards 2030 / 2050 The 2012 Revision. FAO. Rome, Italy. (2012)
- 13. Hall C, Dawson TP, Macdiarmid JI et al. The impact of population growth and climate change on food security in Africa: looking ahead to 2050. Int. J. Agric. Sustain. 15 (2), 124-135 (2017).

- 14. FAO. Climate-Smart Agriculture: Policies, Practices and Financing for Food Security, Adaptation and Mitigation [Internet]. (2010). Available from: http://www.fao.org/doc rep/013/i1881e/i1881e00.htm
- 15. Carter S, Herold M, Rufino MC, Neumann K, Kooistra L, Verchot L. Mitigation of agricultural emissions in the tropics: comparing forest land-sparing options at the national level. Biogeosciences 12(15), 4809-4825 (2015).
- 16. Richards M, Bruun TB, Campbell BM et al. How countries plan to address agricultural adaptation and mitigation. An analysis of Intended Nationally Determined Contributions. CCAFS Info Note (2015).
- 17. FAO. Trade Reforms and Food Security: Conceptualizing the Linkages [Internet]. (2003). Available from: http:// www.fao.org/docrep/005/y4671e/y4671e00.htm#Con
- 18. Hosonuma N, Herold M, De Sy V et al. An assessment of deforestation and forest degradation drivers in developing countries. Environ. Res. Lett.. 7(4), 44009 (2012).
- 19. De Sy V, Herold M, Achard F et al. Land use patterns and related carbon losses following deforestation in South America. Environ. Res. Lett. 10(12), 124004 (2015).
- 20. Kissinger G, Herold M, De Sy V. Drivers of deforestation and forest degradation: a synthesis report for REDD+ policymakers.Lexeme Consulting, Vancouver, Canada (2012).
- 21. Locatelli B, Pavageau C, Pramova E, Di Gregorio M. Integrating climate change mitigation and adaptation in agriculture and forestry: opportunities and trade-offs. Wiley Interdiscip. Rev. Clim. Chang., 6(6), 585–598 (2015).
- 22. Salvini G, Herold M, De Sy V, Kissinger G, Brockhaus M, Skutsch M. How countries link REDD+ interventions to drivers in their readiness plans: implications for monitoring systems. Environ. Res. Lett. 9(7), 74004 (2014).
- 23. Mandemaker M, Bakker M, Stoorvogel J. The role of governance in agricultural expansion and intensification: a global study of arable agriculture. Ecol. Soc. 16(2), art8 (2011).
- 24. Strassburg BBN, Latawiec AE, Creed A et al. Biophysical suitability, economic pressure and land-cover change: a global probabilistic approach and insights for REDD+. Sustain. Sci. 9(2), 129-141 (2013).
- 25. Kastner T, Rivas MJI, Koch W, Nonhebel S. Global changes in diets and the consequences for land requirements for food. Proc. Natl. Acad. Sci. U. S. A. 109(18), 6868-6872 (2012).
- 26. Rudel TK, Schneider L, Uriarte M et al. Agricultural intensification and changes in cultivated areas, 1970-2005. Proc. Natl. Acad. Sci. U. S. A. 106(49), 20675-20680 (2009).
- 27. Minang PA, Van Noordwijk M, Freeman OE, Mbow C, De Leeuw J, Catacutan D. Climate-Smart Landscapes: Multifunctionality in Practice. World Agrforestry Centre (ICRAF), Nairobi, Kenya (2015).
- 28. Ellison D, Morris CE, Locatelli B et al. Trees, forests and water: cool insights for a hot world. Glob. Environ. Chang. 43, 51-61 (2017).
- 29. Vignola R, Locatelli B, Martinez C, Imbach P. Ecosystembased adaptation to climate change: what role for policy-makers, society and scientists? Mitigat. Adapt. Strat. Glob. Chang. 14, 691-696 (2009).
- 30. Turner W, Oppenheimer M, Wilcove D. A force to fight global warming. Nature 462, 278-279 (2009).
- 31. Alkama R, Cescatti A. Biophysical climate impacts of recent changes in global forest cover. Science 351(6273), 600-604 (2016).
- 32. van Noordwijk M, Bayala J, Hairiah K et al. Agroforestry solutions for buffering climate variability and adapting to change. In: Climate Change Impact and Adaptation in

- Agricultural Systems. CAB-International, Wallingford, UK, 216-232 (2014).
- 33. Lipper L, Thornton P, Campbell BM et al. Climate-smart agriculture for food security. Nat. Clim. Chang. 4(December), 1068-1072 (2014).
- 34. Foley JA, Ramankutty N, Brauman KA et al. Solutions for a cultivated planet. Nature 478(20 October), 337–342 (2011).
- 35. Sa JCM, Lal R, Cerri CC, Lorenz K, Hungria M, Carvalho PCF. Low-carbon agriculture in South America to mitigate global climate change and advance food security. Environ. Int..98, 102-112 (2017).
- 36. Climate Action. Brazil's programme on low carbon agriculture [Internet]. (2012). Available from: http://www.cli mateactionprogramme.org/news/brazil s_low_carbon_agriculture_programme
- 37. Carlson KM, Gerber JS, Mueller ND et al. Greenhouse gas emissions intensity of global croplands. Nat. Clim. Chang. 7(January), 63–68 (2016).
- 38. van Noordwijk M, Khasanah N, Dewi S. Can intensification reduce emission intensity of biofuel through optimized fertilizer use? Theory and the case of oil palm in Indonesia. GCB Bioenergy 9, 940-952 (2016).
- 39. Soliman T, Lim FKS, Lee JSH, Carrasco LR. Closing oil palm yield gaps among Indonesian smallholders through industry schemes, pruning, weeding and improved seeds. R. Soc. Open Sci. 3, 1-9 (2016).
- 40. Cohn A, Bowman M, Zilberman D, O'Neill K. The Viability of Cattle Ranching Intensification in Brazil as a Strategy to Spare Land and Mitigate Greenhouse Gas Emissions [Internet]. CCAFS. Copenhagen, Denmark (2011). Available from: https://ccafs.cgiar.org/
- 41. Busch J, Ferretti-Gallon K. What drives deforestation and what stops it? A meta-analysis. Rev. Environ. Econ. Policy. 11(1), 3-23 (2017).
- 42. Lal R. Feeding 11 billion on 0.5 billion hectare of area under cereal crops. Food Energy Secur. 5(4), 239-251
- 43. UNCCD. The Global Land Outlook.Secretariat of the United Nations Convention to Combat Desertification. Bonn, Germany.(2017).
- 44. Minasny B, Malone BP, Mcbratney AB et al. Soil carbon 4 per mille. Geoderma 292, 59-86 (2017).
- 45. Paustian K, Lehmann J, Ogle S, Reay D, Robertson GP, Smith P. Climate-smart soils. Nature 532(7597), 49-57
- 46. Lusiana B, van Noordwijk M, Cadisch G. Land sparing or sharing? Exploring livestock fodder options in combination with land use zoning and consequences for livelihoods and net carbon stocks using the FALLOW model. Agric. Ecosyst. Environ. 159, 145-160 (2012).
- 47. deB Richter D, Houghton RA. Gross CO2 fluxes from landuse change: implications for reducing global emissions and increasing sinks. Carbon Manag. 2(1), 41-47 (2014).
- 48. Zomer RJ, Neufeldt H, Xu J et al. Global tree cover and biomass carbon on agricultural land: the contribution of agroforestry to global and national carbon budgets. Sci. Rep. 6, 29987 (2016).
- 49. Wilkes A, Tennigkeit T, Solymosi K. National Integrated Mitigation Planning in Agriculture: A Review Paper. FAO. Rome, Italy.
- 50. Rosenstock TS, Tully KL, Arias-Navarro C, Neufeldt H, Butterbach-Bahl K, Verchot LV. Agroforestry with N2-fixing trees: sustainable development's friend or foe? Curr. Opin. Environ. Sustain. 6(February), 15-21 (2014).
- 51. Mbow C, Smith P, Skole D, Duguma L, Bustamante M. Achieving mitigation and adaptation to climate change

- through sustainable agroforestry practices in africa. Curr. Opin. Environ. Sustain. 6(1), 8-14 (2014).
- 52. 4 per 1000 Initiative. 4‰ Initiative: soils for food security and climate [Internet]. 4 pour 1000. (2016). Available from: http://4p1000.org/
- 53. Bos JFFP, Hein FM, Verhagen J, Van Ittersum MK. Tradeoffs in soil fertility management on arable farms. Agric. Syst. 157, 292–302 (2017).
- 54. de Vries W. Soil carbon 4 per mille: a good initiative but let's manage not only the soil but also the expectations. Comment on Minasny et al. (2017) Geoderma 292: 59-86. Geoderma 309, 111-112 (2018).
- 55. Giller KE, Andersson JA, Corbeels M et al. Beyond conservation agriculture. Front. Plant Sci. 6(October), Art 870 (2015).
- 56. Valin H, Havlík P, Mosnier A, Herrero M, Schmid E, Obersteiner M. Agricultural productivity and greenhouse gas emissions: trade-offs or synergies between mitigation and food security? Environ. Res. Lett. 8(3), 35019 (2013).
- 57. Havlík P, Valin H, Herrero M et al. Climate change mitigation through livestock system transitions. Proc. Natl. Acad. Sci. U. S. A. 111(10), 3709-3714 (2014).
- 58. Descheemaeker K, Oosting SJ, Tui SH-K, Masikati P, Falconnier GN, Giller KE. Climate change adaptation and mitigation in smallholder crop-livestock systems in sub-Saharan Africa: a call for integrated impact assessments. Reg. Environ. Chang. 16, 2331 (2016).
- 59. Engel S, Muller A. Payments for environmental services to promote "climate-smart agriculture"? Potential and challenges. Agric. Econ. 47, 173–184 (2016).
- 60. Turnhout E, Gupta A, Weatherley-Singh J et al. Envisioning REDD+ in a post-Paris era: between evolving expectations and current practice. Wiley Interdiscip. Rev. Clim. Chang. 8(1), 1-13 (2016).
- 61. Reidsma P, Bakker MM, Kanellopoulos A et al. Sustainable agricultural development in a rural area in the Netherlands? Assessing impacts of climate and socio-economic change at farm and landscape level. Agric. Syst. 141, 160-173 (2015).
- 62. Salvini G, Ligtenberg A, van Paassen A, Bregt AK, Avitabile V, Herold M. REDD+ and climate smart agriculture in landscapes: a case study in Vietnam using companion modeling. J. Environ. Manage. 172, 58-70 (2016).
- 63. Villamor GB, van Noordwijk M. Social role-play games Vs individual perceptions of conservation PES agreements for maintaining rubber agroforests in Jambi (Sumatra), Indonesia. Ecol. Soc. 16(3), 27 (2011).
- Diniz FH, Kok K, Hoogstra-Klein MA, Arts B. Mapping future changes in livelihood security and environmental sustainability based on perceptions of small farmers in the Brazilian Amazon. Ecol. Soc. 20(2), art26 (2015).
- 65. de Sassi C, Joseph S, Bos AB, Duchelle AE, Ravikumar A, Herold M. Towards integrated monitoring of REDD+. Curr. Opin. Environ. Sustain. 14, 93-100 (2015).
- 66. Pratihast A, Herold M, De Sy V, Mudiyarso D, Skutsch M. Linking community-based and national REDD+ monitoring: a review of the potential. Carbon Manag. 4(1), 91-104 (2013).
- 67. Bastos Lima MG, Braña-Varela J, Kleymann H, Carter S. Contribution of forests and land use to closing the gigatonne emissions gap by 2020. WWF, Gland Switzerland (2015).
- 68. Roman-Cuesta RM, Rufino M, Herold M et al. Hotspots of gross emissions from the land use sector: patterns, uncertainties, and leading emission sources for the period 2000-2005 in the tropics. Biogeosciences. 13(14), 4253-4269 (2016).



- 69. Rosenstock TS, Rufino MC, Butterbach-Bahl K, Wollenberg E. Toward a protocol for quantifying the greenhouse gas balance and identifying mitigation options in smallholder farming systems. Environ. Res. Lett. 8 (2), 21003 (2013).
- 70. Dinesh D, Vermeulen S, Bacudo I et al. Options for agriculture at Marrakech climate talks: messages for SBSTA 45 agriculture negotiators. CCAFS Rep. 16. Copenhagen, Denmark. (2016).
- 71. Chandra A, McNamara KE, Dargusch P et al. Resolving the UNFCCC divide on climate-smart agriculture. Carbon Manag. 7(5-6), 295-299 (2016).
- 72. Den Besten JW, Arts B, Verkooijen P. The evolution of REDD+: an analysis of discursive-institutional dynamics. Environ. Sci. Policy 35, 40-48 (2014).
- 73. Gutiérrez-Vélez VH, DeFries R, Pinedo-Vásquez M et al. High-yield oil palm expansion spares land at the expense of forests in the Peruvian Amazon. Environ. Res. Lett. 6(4), 44029 (2011).
- 74. Singh VP, Sinha RB, Nayak D, Neufeldt H, Van Noordwijk M, Rizvi J. The National Agroforestry Policy of India: Experiential Learning in Development and Delivery Phases [Internet]. Working Paper 240. World Agroforestry Centre (ICRAF), New Delhi (India). (2016) Available from: http:// www.worldagroforestry.org/downloads/Publications/ PDFS/WP16143.pdf
- 75. Grassi G, House J, Dentener F, Federici S, den Elzen M, Penman J. The key role of forests in meeting climate targets requires science for credible mitigation. Nat. Clim. Chang. 7(February), 220–225 (2017).
- 76. DeFries RS, Rudel T, Uriarte M, Hansen M. Deforestation driven by urban population growth and agricultural trade in the twenty-first century. Nat. Geosci. 3(3), 178-181 (2010).
- 77. Brown S, Zarin D. What does zero deforestation mean? Science 342(6160), 805-807 (2013).
- 78. Nepstad D, Irawan S, Bezerra T et al. More food, more forests, fewer emissions, better livelihoods: linking REDD+, sustainable supply chains and domestic policy in Brazil, Indonesia and Colombia. Carbon Manag. 4(6), 639-658 (2013).
- 79. Stickler C, Bezerra T, Nepstad D. Global rules for sustainable farming: a comparison of social and environmental safeguards for REDD+ and principles & criteria for commodity roundtables. The RT-REDD Consortium (2012).
- 80. WWF. Jurisdictional approaches to zero deforestation commodities. WWF Discussion paper (2016).
- 81. Earth Innovation Institute. Forests, farms & finance initiative [Internet]. (2016). Available from: http://earthinnova tion.org/our-work/global/forests-farms-finance-initiative/
- 82. Consumer Goods Forum. Acting individually: production protection [Internet]. (2015). Available from: http:// tfa2020.org/wp-content/uploads/2015/12/01122015-__Produce-Protect-CGF-statement.pdf

- 83. Earth Innovation Institute. Making corporate deforestation pledges work [Internet]. (2016). Available from: http://earthinnovation.org/wp-content/uploads/2014/ 09/CDZ-report_online-1.pdf
- 84. Smith P, Haberl H, Popp A et al. How much land-based greenhouse gas mitigation can be achieved without compromising food security and environmental goals? Glob. Chang. Biol. 19(8), 2285-2302 (2013).
- 85. Van Kernebeek HRJ, Oosting SJ, Van Ittersum MK, Bikker P, De Boer IJM. Saving land to feed a growing population: consequences for consumption of crop and livestock products. Int. J. Life Cycle Assess. 21(5), 677-687 (2016).
- 86. Erb K-H, Lauk C, Kastner T, Mayer A, Theurl MC, Haberl H. Exploring the biophysical option space for feeding the world without deforestation. Nat. Commun. 7, 1-7 (2016).
- 87. Derkx B, Glasbergen P. Elaborating global private metagovernance: an inventory in the realm of voluntary sustainability standards. Glob. Environ. Chang. 27(1), 41–50 (2014).
- 88. Mithöfer D, van Noordwijk M, Leimona B, Cerutti P. Certify and shift blame, or resolve issues? Environmentally and socially responsible global trade and production of timber and tree crop commodities. J. Biodivers. Sci. Ecosyst. Serv. Manag. 13(1), 72-95 (2016).
- 89. CDP. Carbon Disclosure Project [Internet]. (2017). Available from: www.cdp.net
- 90. Doda B, Gennaioli C, Gouldson A, Grover D, Sullivan R. Are corporate carbon management practices reducing corporate carbon emissions? Corp. Soc. Responsib. Environ. Manag. 23(5), 257-270 (2016).
- 91. De Sy V, Herold M, Martius C et al. Enhancing Transparency in the Land-Use Sector: Exploring the Role of Independent Monitoring Approaches [Internet]. CIFOR, Bogor, Indonesia. (2016) Available from: http://www.cifor.org/ library/6256/enhancing-transparency-in-the-land-use-sec tor-exploring-the-role-of-independent-monitoringapproaches/
- 92. World Resources Institute. Global Forest Watch [Internet]. (2017). Available from: http://www.globalforestwatch.org/
- 93. FAO. OpenForis [Internet]. (2017). Available from: http:// www.openforis.org/
- 94. Janssen SJC, Porter CH, Moore AD et al. Towards a new generation of agricultural system data, models and knowledge products: information and communication technology. Agric. Syst. 155, 200-212 (2017).
- 95. UNFCCC. Report of the Conference of the Parties on Its Eighth Session: 17/CP.8 [Internet]. New Delhi, India (2003) Available from: https://unfccc.int/files/meetings/work shops/other_meetings/application/pdf/dec17-cp.pdf
- 96. Gupta A, Mason M. Disclosing or obscuring? The politics of transparency in global climate governance. Curr. Opin. Environ. Sustain. 18, 82-90 (2016).